

Super-resolution with the Nanoimager what is the most suitable imaging technique for you?

	dSTORM Direct Stochastic Optical Reconstruction Microscopy	PALM PhotoActivated Localization Microscopy	SPT Single-Particle Tracking	smFRET Single-molecule Förster Resonance Energy Transfer
Resolution achieved	20nm	20nm	N/A	0.3-10nm
# of colors	Up to 4 (2 simultaneously)	Up to 4 (2 simultaneously)	Up to 4 (2 simultaneously)	2 FRET pairs at once
Live or fixed samples?	Mostly fixed	Both	Live (in solution or cells)	Both
Best for	Quantification and localization of single moleculesVery high resolution	Counting and tracking of single molecules in fixed & live samples Lower laser powers than for dSTORM	Tracking single particles and molecules Quantitative data (i.e. diffusion coeff., sizing and counting)	Real-time measurement of interor intramolecular interactions Individual traces and population averages
Added value	TIRF/HILO for a better SNR Tip Line Line Line Line Line Line Line Line	TIRF/HILO for a better SNR TIRF/HILO for a better SNR Astigmatic 3D Whole body heating for live imaging	TIRF/HILO for a better SNR TIRF/HILO for a better SNR Astigmatic 3D Whole body heating for live imaging Microfluidics compatibility	Use of Alex (Alternating Laser EXcitation) mode for stoichiometry of molecules
Nanoimager	COM COM COM			ON ON ON
Data analysis	Q' Q' Q' Q'	Q ' Q ' Q ' Q '	Q ' Q ' Q ' Q '	Q Q Q
Data analysis tools on Nanoimager	O Python	python •	python of	python:
Pro tips	Choose blinking dyes and special buffersVery clean coverslips	Photoactivable and photoconvertible fluorophores Very clean coverslips	Choose bright and long-emitting fluorophoresVery clean coverslips	Donor emission to overlap with acceptor's absorption spectrum Exceptionally clean coverslips
Additional notes	Live imaging not feasible due to high laser power and non-physiological buffers Longer acquisition times	Limited availability of fluorophores Acquisition can be time-consuming (depending on number of molecules)	Tricky to use in densely labeled population Fluorophores need to be sparse	Only works over short distances (up to 10nm) Compatible FRET pairs required





00:00 Real-time rendering







Dedicated smFRET analysis tool

For the best flourophores and imaging guides visit: www.oni.bio/resources/oniversity



